

Another point in homological algebra: Duality for discontinuous group actions

Fritz Grunewald and Wilhelm Singhof

Heinrich-Heine-Universität Düsseldorf

e-mail: fritz@math.uni-duesseldorf.de

January 16, 2009

Abstract

We consider discontinuous operations of a group G on a contractible n -dimensional manifold X . Let E be a finite dimensional representation of G over a field k of characteristics 0. Let $\mathcal{E}$ be the sheaf on the quotient space $Y = G \backslash X$ associated to E . Let $H_!^\bullet(Y; \mathcal{E})$ be the image in $H^\bullet(Y; \mathcal{E})$ of the cohomology with compact support. In the cases where both $H_!^\bullet(Y; \mathcal{E})$ and $H_!^\bullet(Y; \mathcal{E}^*)$ ($\mathcal{E}^*$ being the sheaf associated to the representation dual to E) are finite dimensional, we establish a non-degenerate duality between $H_!^m(Y; \mathcal{E})$ and $H_!^{n-m}(Y; \mathcal{E}^*)$. We also show that this duality is compatible with Hecke operators.

AMS classification: 20J05, 11F06, 14L30, 16E99, 20G10

Contents

1	Introduction	1
2	Poincaré duality	2
3	Hecke operators	7

1 Introduction

We consider discontinuous operations of a group G on a contractible n -dimensional manifold X . Let E be a finite dimensional representation of

G over a field k of characteristics 0, say. Associated to E is a sheaf $\mathcal{E}$ on the quotient space $Y = G \setminus X$, and a theorem of Grothendieck [G] says that there is an isomorphism

$$H^\bullet(G; E) \cong H^\bullet(Y; \mathcal{E}) . \quad (1)$$

This is a result which is basic for the cohomology theory of groups. In $H^\bullet(Y; \mathcal{E})$, we have the canonical subgroup $H_!^\bullet(Y; \mathcal{E})$ which is the image of the cohomology with compact supports. In Section 2, we show that there is a non-degenerate pairing

$$B : H_!^m(Y; \mathcal{E}) \otimes H_!^{n-m}(Y; \mathcal{E}^*) \rightarrow k . \quad (2)$$

This of course gives rise to isomorphisms

$$H_!^m(Y; \mathcal{E})^* \cong H_!^{n-m}(Y; \mathcal{E}^*) \quad (3)$$

where $\mathcal{E}^*$ is the sheaf associated to the dual representation E^* and where $H_!^m(Y; \mathcal{E})^*$ is the dual of $H_!^m(Y; \mathcal{E})$.

Special instances of this form of Poincaré duality were frequently used. In full generality, we derive it by using Grothendieck's arguments together with Poincaré-Verdier duality. In fact we feel that our paper is another (perhaps missing) chapter in the work [G] of Grothendieck.

In Section 3, we show that the above duality isomorphism (3) is compatible with Hecke operators.

2 Poincaré duality

Let X be a connected, orientable, n -dimensional manifold and let G be a discrete group operating effectively and smoothly on X preserving the orientation. We assume that this operation is *discontinuous*, i.e. the following two conditions are satisfied:

- For each $x \in X$, the stabilizer G_x is finite.
- Each x admits a neighborhood V such that

$$\gamma \cdot V \cap V = \emptyset \quad \text{for all } \gamma \in G \setminus G_x .$$

We write $Y := G \setminus X$ and denote by $p : X \rightarrow Y$ the natural projection. Let X_{reg} be the open subset of X consisting of all points with trivial stabilizer.

Since the smooth operation of a finite group looks, in the neighborhood of a fixed point of the group, like a linear operation [B1], we have the following two properties:

Fact 2.1 $X \setminus X_{\text{reg}}$ has at least codimension 2 in X .

Fact 2.2 Every point of Y has an open neighborhood U such that there exists an open subset V of X diffeomorphic to $\mathbb{R}^n$ and a finite subgroup H of G such that $H \cdot V = V$ and that p induces a homeomorphism from $H \setminus V$ onto U .

Since paths in quotients by finite groups can be lifted ([B1], Th. 6.2, p. 91), we conclude from 2.2:

Fact 2.3 Let A be a pathwise connected subset of Y and let A_0 be a path component of $p^{-1}(A)$. Then:

- a) $p^{-1}(A) = G \cdot A_0$.
- b) For each $\gamma \in G$, either $\gamma \cdot A_0 = A_0$ or $\gamma \cdot A_0 \cap A_0 = \emptyset$.

Let k be a field with the following property: For each $x \in X$, the characteristics of k is prime to the order of G_x .

With the terminology of [B2], Def. V.9.1 and using [B2], Th. II.19.2 and [B2], Th. V.19.3, we then have:

Fact 2.4 Y is an orientable n -dimensional homology manifold over k .

By a representation of G , we mean a finite dimensional vector space E over k with a linear operation of G on it. A sheaf will always be a sheaf of k -vector spaces.

We consider the following functor $\mathcal{S}$ from the category of representations of G to the category of sheaves on Y : For a representation E , the sheaf $\mathcal{E} := \mathcal{S}(E)$ is given by

$$\Gamma(U; \mathcal{E}) := \{\text{locally constant } G\text{-equivariant maps } p^{-1}(U) \rightarrow E\}$$

for open subsets U of Y . By 2.3, the stalks of $\mathcal{E}$ are given by

$$\mathcal{E}_y = \{G\text{-equivariant maps } p^{-1}(y) \rightarrow E\}.$$

The constant sheaf on X with stalks E , denoted by E_X , is a special instance of a G -sheaf on X . For any G -sheaf $\mathcal{F}$ on X , the direct image sheaf $p_*(\mathcal{F})$ is a G -sheaf on Y , where G operates trivially on Y . Let $p_*^G(\mathcal{F})$ be the subsheaf of $p_*(\mathcal{F})$ consisting of those elements which are fixed under G . With this notation, we have

$$\mathcal{S}(E) = p_*^G(E_X).$$

More generally, following [G] (5.2.4), we introduce the sheaves

$$\mathcal{H}^m(G; \mathcal{F}) := R^m p_*^G(\mathcal{F})$$

on Y so that $\mathcal{S}(E) = \mathcal{H}^0(G; E_X)$.

We denote by $\Gamma_X^G(\mathcal{F})$ the G -invariant sections of $\mathcal{F}$ and put as in [G] (5.2.3):

$$H^m(X; G, \mathcal{F}) := R^m \Gamma_X^G(\mathcal{F}) .$$

Moreover, let $\Gamma_{X,c}^G(\mathcal{F})$ denote the elements in $\text{Gamma}_X^G(\mathcal{F})$ with compact support and put as in [G] (5.7.2):

$$H_c^m(X; G, \mathcal{F}) := R^m \Gamma_{X,c}^G(\mathcal{F}) .$$

By [G], Th. 5.2.1 there are two spectral sequences

$$(H^p(Y; \mathcal{H}^q(G; \mathcal{F})))_{p,q} \Rightarrow H^\bullet(X; G, \mathcal{F}) , \quad (4)$$

$$(H^p(G; H^q(X; \mathcal{F})))_{p,q} \Rightarrow H^\bullet(X; G, \mathcal{F}) . \quad (5)$$

For a *finite* group G , there are, by [G] (5.7.4) and (5.7.7), analogs for the cohomology with compact supports.

$$(H_c^p(Y; \mathcal{H}^q(G; \mathcal{F})))_{p,q} \Rightarrow H_c^\bullet(X; G, \mathcal{F}) , \quad (6)$$

$$(H^p(G; H_c^q(X; \mathcal{F})))_{p,q} \Rightarrow H_c^\bullet(X; G, \mathcal{F}) . \quad (7)$$

From the spectral sequences (3) and (4), Grothendieck deduces the basic relation (1). Similarly, from (5) and (6) we get:

Fact 2.5 *If the group G is finite and if $\text{char } k$ is prime to the order of G , we have*

$$H_c^m(Y; \mathcal{E}) \cong H_c^m(X; E)^G .$$

For two sheaves $\mathcal{F}, \mathcal{G}$ on Y and $p \geq 0$, we have the sheaf $\mathcal{E}xt^p(\mathcal{F}, \mathcal{G})$. It is the sheafification of the presheaf

$$U \mapsto \text{Ext}^p(\mathcal{F}|_U, \mathcal{G}|_U) .$$

We write $\mathcal{H}om$ instead of $\mathcal{E}xt^0$. Observe that the presheaf

$$U \mapsto \text{Hom}(\mathcal{F}|_U, \mathcal{G}|_U)$$

is a sheaf, hence $\mathcal{H}om(\mathcal{F}, \mathcal{G})(U) = \text{Hom}(\mathcal{F}|_U, \mathcal{G}|_U)$.

Lemma 2.6 *For two representations E, F of G and their associated sheaves $\mathcal{E}, \mathcal{F}$, there is a canonical isomorphism*

$$\mathcal{S}(\mathrm{Hom}_k(E, F)) \cong \mathcal{H}om(\mathcal{E}, \mathcal{F}).$$

Proof. Writing $\mathcal{G} := \mathcal{S}(\mathrm{Hom}_k(E, F))$, we have a canonical map

$$\alpha : \mathcal{G} \rightarrow \mathcal{H}om(\mathcal{E}, \mathcal{F}).$$

The non-trivial point is to show that

$$\alpha_y : \mathcal{G}_y \rightarrow \mathcal{H}om(\mathcal{E}, \mathcal{F})_y$$

is surjective for all $y \in Y$. We have

$$\mathcal{G}_y = \{G\text{-equivariant maps } \varphi : p^{-1}(y) \rightarrow \mathrm{Hom}_k(E, F)\}.$$

Fixing a point $x \in p^{-1}(y)$, the evaluation $\varphi \mapsto \varphi_x$ gives an isomorphism

$$\mathcal{G}_y \cong \mathrm{Hom}_{k_{G_x}}(E, F).$$

Given an element $\delta \in \mathcal{H}om(\mathcal{E}, \mathcal{F})_y$, we choose an open neighborhood U of y in Y which is so small that each component of $p^{-1}(U)$ contains exactly one point of $p^{-1}(y)$ and that δ can be represented by an element $\psi \in \mathrm{Hom}(\mathcal{E}|_U, \mathcal{F}|_U)$. Let U_0 be the component of $p^{-1}(U)$ containing x . Let $\xi \in U_0 \cap X_{\mathrm{reg}}$ and $\eta := p(\xi)$. Evaluation in ξ gives isomorphisms $\mathcal{E}_\eta \cong E$ and $\mathcal{F}_\eta \cong F$. Hence the homomorphism $\psi_\eta : \mathcal{E}_\eta \rightarrow \mathcal{F}_\eta$ gives a k -homomorphism $\varphi : E \rightarrow F$. Using 2.1, it is easy to see that φ doesn't depend on the choice of ξ . Moreover, this implies that $\varphi \in \mathrm{Hom}_{k_{G_x}}(E, F) \cong \mathcal{G}_y$. It is clear that $\alpha_y(\varphi) = \delta$. $\square$

Given a representation E of G , we denote by $E^* := \mathrm{Hom}_k(E, k)$ the dual representation and write $\mathcal{E}^* := \mathcal{S}(E^*)$.

Corollary 2.7 $\mathcal{H}om(\mathcal{E}, \mathcal{F}) \cong \mathcal{H}om(\mathcal{F}^*, \mathcal{E}^*)$.

Because of 2.4, we can apply Poincaré-Verdier duality (see e.g. [GM], (III. 70)) and obtain:

$$\mathrm{Hom}_k(H_c^m(Y; \mathcal{E}), k) \cong \mathrm{Ext}^{n-m}(\mathcal{E}, k_Y). \quad (8)$$

Lemma 2.8 $\mathrm{Ext}^p(\mathcal{E}, k_Y) = 0$ for $p > 0$.

Proof. This is a local property. By 2.2 it suffices to assume that X is diffeomorphic to $\mathbb{R}^n$ and that G is finite. We have

$$\begin{aligned} \mathrm{Ext}^p(\mathcal{E}, k_Y) &\cong \mathrm{Hom}_k(H_c^{n-p}(Y; \mathcal{E}), k) && \text{by (7)} \\ &\cong \mathrm{Hom}_k(H_c^{n-p}(X; E)^G, k) && \text{by 2.5.} \end{aligned}$$

Since $X \approx \mathbb{R}^n$, we have $H_c^{n-p}(X; E) = 0$ for $p > 0$. $\square$

Lemma 2.9 *For each $p \geq 0$, there is a canonical isomorphism*

$$\mathrm{Ext}^p(\mathcal{E}, k_Y) \cong H^p(Y; \mathcal{E}^*) .$$

Proof. We use the spectral sequence ([G], Th. 4.2.1)

$$(H^p(Y; \mathcal{E}xt^q(\mathcal{E}, k_Y)))_{p,q} \Rightarrow \mathrm{Ext}^\bullet(\mathcal{E}, k_Y) .$$

By 2.8, we obtain $\mathrm{Ext}^p(\mathcal{E}, k_Y) \cong H^p(Y; \mathcal{H}om(\mathcal{E}, k_Y))$. Then apply 2.7. $\square$

There is a canonical homomorphism of sheaves

$$\mu : \mathcal{E} \otimes \mathcal{E}^* \rightarrow k_Y$$

which induces

$$\mu_* : H_c^n(Y; \mathcal{E} \otimes \mathcal{E}^*) \rightarrow H_c^n(Y; k) .$$

Composing with the cup-product

$$\cup : H_c^m(Y; \mathcal{E}) \otimes H^{n-m}(Y; \mathcal{E}^*) \rightarrow H_c^n(Y; \mathcal{E} \otimes \mathcal{E}^*)$$

and with

$$\int : H_c^n(Y; k) \rightarrow k ,$$

we obtain for each $m \geq 0$ a homomorphism

$$B : H_c^m(Y; \mathcal{E}) \otimes H^{n-m}(Y; \mathcal{E}^*) \rightarrow k .$$

Combining Poincaré-Verdier duality (7) and in particular the remark in [GM] following the formula (III.70) with 2.9, we obtain

Lemma 2.10 *The pairing B is non-singular for each $m \geq 0$.* $\square$

For a sheaf $\mathcal{F}$ on Y , let us write

$$H_i^\bullet(Y; \mathcal{F}) := \mathrm{im}(H_c^\bullet(Y; \mathcal{F}) \rightarrow H^\bullet(Y; \mathcal{F})) .$$

The properties of the cup-product, in particular its anti-commutativity, show that 2.10 implies the following duality for $H_i^\bullet$:

Theorem 2.11 *Let X be an orientable, connected, n -dimensional manifold and let G be a group operating smoothly and discontinuously on X preserving the orientation. Let k be a field such that for each $x \in X$, the characteristics of k is prime to the order of the stabilizer G_x . Let E be a representation of G on a finite-dimensional k -vector space and let E^* be the dual representation. We denote by $\mathcal{E}$ the sheaf on $Y = G \backslash X$ associated to E and by $\mathcal{E}^*$ the sheaf associated to E^* and assume that all cohomology groups $H^m(Y; \mathcal{E})$ and $H^m(Y; \mathcal{E}^*)$ are finite dimensional k -vector spaces. Then B induces for each $m \geq 0$ a non-singular pairing of finite dimensional k -vector spaces*

$$B : H_i^m(Y; \mathcal{E}) \otimes H_i^{n-m}(Y; \mathcal{E}^*) \rightarrow k . \quad (9)$$

3 Hecke operators

Let us now assume in addition that there is a group $\mathfrak{G}$ with

$$G \subseteq \mathfrak{G} \subseteq \text{Diff } X$$

and that E is actually a representation of $\mathfrak{G}$. Let $g \in \mathfrak{G}$ be an element in the commensurator of G , that is, we assume that the two groups

$$\begin{aligned} \Delta' &:= gGg^{-1} \cap G, \\ \Delta'' &:= G \cap g^{-1}Gg = g^{-1}\Delta'g \end{aligned}$$

are of finite index in G . Consider the following diagram:

$$\begin{array}{ccccc} X & \xrightarrow{p'} & \Delta' \setminus X & \xrightarrow{\varphi} & \Delta'' \setminus X & \xleftarrow{p''} & X \\ & \searrow p & \downarrow \pi' & & \downarrow \pi'' & \swarrow p & \\ & & G \setminus X & & G \setminus X & & \end{array} \quad (10)$$

All the arrows except φ are natural projections; $\varphi = \varphi_g$ is the diffeomorphism given by

$$\varphi(\Delta'x) := \Delta''g^{-1}x = g^{-1}\Delta'x .$$

As we shall explain, we obtain from (8) homomorphisms

$$\begin{array}{ccc} H^\bullet(\Delta' \setminus X; \mathcal{E}') & \xleftarrow{\varphi^*} & H^\bullet(\Delta'' \setminus X; \mathcal{E}'') \\ \downarrow \tau & & \uparrow \pi''^* \\ H^\bullet(G \setminus X; \mathcal{E}) & & H^\bullet(G \setminus X; \mathcal{E}) \end{array} \quad (11)$$

Here $\mathcal{E}$ is the sheaf introduced in section 2, and $\mathcal{E}'$ (resp. $\mathcal{E}''$) are the analogous sheaves on $\Delta' \setminus X$ (resp. $\Delta'' \setminus X$).

Definition. The *Hecke operator* $T(g)$ is the endomorphism of $H^\bullet(G \setminus X; \mathcal{E})$ given by

$$T(g) := \tau \circ \varphi^* \circ \pi''^* .$$

Let us now explain the three homomorphisms τ , φ^* , and π''^* in (9): We have a canonical inclusion $i' : \mathcal{E} \hookrightarrow \pi'_* \mathcal{E}'$. We obtain induced homomorphisms

$$\pi'^{\dagger} : H^\bullet(G \setminus X; \pi'_* \mathcal{E}') \rightarrow H^\bullet(\Delta' \setminus X; \mathcal{E}')$$

which is an isomorphism by the Vietoris mapping theorem ([B2], Th. II. 11.1) and

$$\pi'^* = \pi'^\dagger \circ i'_* : H^\bullet(G \setminus X; \mathcal{E}) \rightarrow H^\bullet(\Delta' \setminus X; \mathcal{E}') .$$

Similarly for $\mathcal{E}''$ instead of $\mathcal{E}'$. This explains π''^* .

Next, there is a sheaf homomorphism

$$\sigma : \pi'_* \mathcal{E}' \rightarrow \mathcal{E}$$

defined as follows: Choose elements $\gamma_1, \dots, \gamma_m \in G$ with

$$G = \coprod_{i=1}^m \gamma_i \Delta' .$$

For U open in $G \setminus X$, define

$$\begin{aligned} \sigma_U : \Gamma(U; \pi'_* \mathcal{E}') &\rightarrow \Gamma(U; \mathcal{E}) \\ \sigma_U(f)(x) &:= \sum_{i=1}^m \gamma_i \cdot f(\gamma_i^{-1} \cdot x) \end{aligned}$$

for $x \in p^{-1}(U)$. The definition of σ doesn't depend on the choice of the γ_i . We then have a *transfer*

$$\tau := \sigma_* \circ (\pi'^\dagger)^{-1} : H^\bullet(\Delta' \setminus X; \mathcal{E}') \rightarrow H^\bullet(G \setminus X; \mathcal{E}) . \quad (12)$$

Observe that τ is a special case of the transfer introduced in [B2], p.140-141. (Bredon's assumption that G is a finite group is not at all used for the definition of the transfer called $\mu_{G/H}^*$ by him.)

Finally, to explain φ^* , we have to specify a φ -cohomomorphism

$$\Phi : \mathcal{E}'' \rightsquigarrow \mathcal{E}'$$

(using the terminology of [B2], I. 4): For each $z \in \Delta' \setminus X$, we need a homomorphism

$$\Phi_z : \mathcal{E}''_{\varphi(z)} \rightarrow \mathcal{E}'_z .$$

Choosing an element $x \in p'^{-1}(z)$, we have

$$\begin{aligned} \mathcal{E}''_{\varphi(z)} &= \{ \Delta''\text{-equivariant maps } f : g^{-1} \Delta' x \rightarrow E \} , \\ \mathcal{E}'_z &= \{ \Delta'\text{-equivariant maps } \Delta' x \rightarrow E \} . \end{aligned}$$

We put

$$\Phi_z(f) : \delta' x \mapsto g \cdot f(g^{-1} \delta' x) .$$

Observe that this is the place where we actually use that E is a representation of the group $\mathfrak{G}$. In any case, together with the cohomomorphism Φ , we obtain the induced homomorphism φ^* . This completes the definition of the Hecke operator $T(g)$.

The same construction works for cohomology with compact supports. Hence we get also Hecke operators

$$\begin{aligned} T(g) : H_c^\bullet(G \setminus X; \mathcal{E}) &\rightarrow H_c^\bullet(G \setminus X; \mathcal{E}) , \\ T(g) : H_!^\bullet(G \setminus X; \mathcal{E}) &\rightarrow H_!^\bullet(G \setminus X; \mathcal{E}) , \end{aligned}$$

The following properties are easy to verify:

Fact 3.1 *On $H^0(G \setminus X; \mathcal{E}) = E^G$, the operator $T(g)$ is given by*

$$T(g) \cdot e = \sum_{i=1}^m \gamma_i g \cdot e$$

where the γ_i are as above.

Fact 3.2 *If γ, γ' are in G , we have*

$$T(\gamma g \gamma') = T(g) .$$

The transfer τ has the following well-known properties:

- $\tau \circ \pi'^*$ is multiplication by the number $[G : \Delta']$.
- $\tau : H^0(\Delta' \setminus X; k) \rightarrow H^0(G \setminus X; k)$ is the homomorphism $[G : \Delta'] : k \rightarrow k$.
- The following diagram is commutative (use 2.1):

$$\begin{array}{ccc} H_c^n(\Delta' \setminus X; k) & \xrightarrow{\tau} & H_c^n(G \setminus X; k) \\ & \searrow f & \swarrow f \\ & k & \end{array}$$

- $\tau(b \cup \pi'^*(c)) = \tau(b) \cup c$ (cf. [B2], p.405)

From these formulas, we conclude that Hecke operators behave well with respect to the pairing

$$B : H_c^m(Y; \mathcal{E}) \otimes H^{n-m}(Y; \mathcal{E}^*) \rightarrow k .$$

from section 2: We have

$$B(T(g) \cdot u \otimes v) = B(u \otimes T(g^{-1}) \cdot v) . \quad (13)$$

So far in this section, we haven't used any restriction on the characteristics of k ; it is not even necessary that k is a field. Now we see:

Theorem 3.3 *Under the assumptions of the duality theorem 2.11, the pair $(H_I^m(Y; \mathcal{E}), T(g))$ consisting of a vector space and an endomorphism is dual to the pair $(H_I^{n-m}(Y; \mathcal{E}^*), T(g^{-1}))$.*

References

- [B1] Bredon, G.E.: Introduction to Compact Transformation Groups. Academic Press, 1972
- [B2] Bredon, G.E.: Sheaf Theory. Second Edition. Springer GTM 170, 1997
- [GM] Gelfand, S.I., Manin, Yu.I.: Methods of Homological Algebra. Springer, 1996
- [G] Grothendieck, A.: Sur quelques points d'algèbre homologique. Tohoku Math. J. **9** (1957), 119-221